# Skeletal System

# Bone or Osseous Tissue

- Connective tissue in which the matrix is hardened by the deposition of calcium phosphate and other minerals.
- Mineralization or calcification- the hardening of bone.

# Functions

- Support
- Protection
- Movement
- Mineral storage and homeostasis
  - calcium and phosphorus
- Blood production- red bone marrow
- Storage of energy
  - yellow bone marrow contains mostly adipose tissue
- Electrolyte balance-buffers blood against excessive pH changes by absorbing or releasing alkaline salt
- Detoxification- removes heavy metals and other foreign elements from the blood, thus reducing their effects on the nervous and other tissue.

# Types of Bones

Classified into four types according to shape

- Long-
- Short
- Flat
- Irregular

Two types classified according to location

- **Sutural (Wormian) bones**- small bones between joints of cranial bones
- **Sesamoid bones**- small bones located in tendons (knee caps)


SAGITTAL SUTURE
PARIETAL BONE
WORMIAN BONE
LAMBOIDAL SUTURE
OCCIPITAL BONE

# Bone According to Structure

- **Compact bone-** dense osseous tissue
- **Spongy (cancellous) bone-** loosely organized osseous tissue found at the end of long bones and in the middle of most others.



(a)

(b)



(c)



(d)


Crack


Outer compact bone
Suture
Spongy bone
Inner compact bone

**Diploe (**DIP-lo-ee)- the spongy layer of bone in the skull that is surrounded by one plates of compact bone on each side.



Kenneth S. Saladin, ANATOMY AND PHYSIOLOGY: THE UNITY OF FORM AND FUNCTION, Copyright © 1998, The McGraw-Hill Companies, Inc. All rights reserved.



# Flat Bone Anatomy

**Diploe (**DIP-lo-ee)- the spongy layer of bone in the skull that is surrounded by one plates of compact bone on each side.



# Parts of Long Bones

- **Diaphysis**- the shaft or long main cylindrical portion
- **Epiphysis**- the ends
- **Metaphysis**- the region where the diaphysis joins the diaphysis
- **Articular cartilage**- thin layer of hyaline cartilage covering the epiphysis
- **Periosteum**- tough white fibrous membrane around the surface of the bone not covered by the articular cartilage
  - consist of dense, irregular connective tissue, blood vessels, and nerves that pass into the bones
- **medullary cavity**- central part of the bone, also called the marrow cavity because it contains the bone marrow.
- **Endosteum**- the lining of the medullary cavity that consists of osteoprogenitor cells and osteoclasts.
- **Nutrient foramina-** minute hole through which blood vessels enter the bone.


Proximal epiphysis
Spongy bone
Compact bone
Nutrient foramen
Nutrient vessel
Medullary cavity
Diaphysis
Perforating fibers
Periosteum
Endosteum
Epiphyseal line
Distal epiphysis
Articular cartilage
Coxa

# Bone Marrow

- Soft tissue that occupies the medullary cavity
  - red bone marrow- hemopoietic tissue (blood producing)
    - in the child most of the bone marrow is red
  - yellow bone marrow- in adults most of the medullary cavities are replaced with yellow bone marrow.
    - red marrow limited to vertebrae, ribs. sternum, part of the pelvic girdle, and the proximal ends of the humerous and femur
  - gelatinous marrow- by old age most of the yellow marrow is replaced by this reddish jell.

# Periosteum

- Composed of
  - tough outer fibrous layer of collagen
    - some fibers are continuous with the tendons
    - some fibers penetrate into the bone matrix as perforating (Sharpey) fibers
      - anchor the periosteum to the circumferential lamellae
  - inner osteogenic layer of bone-forming cells (osteogenic cells)



# Bone Histology

The bone matrix has two main components :

- **Organic matrix**
  - gives bone its strength.
    - composed of **type I collagen fibers** (about 95%) embedded in an **amorphous ground substance** consisting of:
      - **sulfated glycosaminoglycans** (chondroitin-4-sulfate, chondroitin-6-sulfate, keratan sulfate)
      - various **bone proteins** (bone sialoprotein, osteocalcin)
- **Inorganic salts.**
  - consist of inorganic components (mineral salts) which makes bone hard.
  - the main calcium deposits in the bone matrix are in the form of crystals of **hydroxyapatite** Ca10(PO4)6.(OH)2

# Bone Histology

- Four types of cells
  - **osteoprogenitor (osteogenic) cells** are precursor bone cells that undergo mitosis to become osteoblast.
    - found in the periosteum, endosteum, and canals in bone that contain blood vessels.
  - **osteoblast**- the cells that form bone but do not have the ability to divide by mitosis.
    - Are found on the surface of bone
    - they initially form collagen and other organic compounds to build bone
  - **osteocyte**- are mature bone cells derived from osteoblast that have secreted bone tissue around themselves.
    - They maintain healthy bone
    - do not undergo mitosis
  - **osteoclast**- function in bone destruction (resorption) which is important in the development, growth, maintenance, and repair of bone.
    - Are multicellular cells that secrete enzymes that breakdown bone


Osteoprogenitor cell
(develops into an
osteoblast)
Osteoblast
(forms bone
tissue)
Osteocyte
(maintains
bone tissue)
Osteoclast
(functions in resorption,
the destruction of bone matrix)

# Osteoblast





# Osteoclast



# Osteoblast



# Osteocytes

# Osteoclast



# Osteoblast, Osteocyte, Osteoclast

# Osteoclast and Howship's Lacunae



# Bone Histology (cont.)

- **calcification-** the deposition and the subsequent hardening of the mineral salts in bone
- bones are composed of channels through which blood vessels supply the nutrients
  - these spaces make bone lighter
  - bones can be classified as compact or spongy, depending on size and location of spaces

# Demineralization of the Clavicle

# Bone Histology (cont.)

Compact (dense) bone

- contains few spaces
- forms external layer of all bone and the bulk of the shaft of long bones
- provides strength and protection
- has a concentric structure
  - Each layer called a **lamellae**
- **Outer circumferential lamellae-** a continuous layers of compact bone at the periphery of the diaphysis



# Bone Histology (cont.)

Compact bone (cont)

- nutrient arteries and nerves from the periosteum penetrate compact bone through **perforating (Volkmann's) canals.**
- **central (Haversian) canals**- canals that run lengthwise through the bone and connect with the perforating canals.
- **Concentric lamellae**- rings of hard, calcified matrix around the central canals.
- **Lacunae**- small spaces between the lamellae
- **Canaliculi**- minute canals that project outward in all directions from the lucunae.
  - Contain slender processes of osteocytes
  - forms system of interconnected canals for nutrients and oxygen to reach osteocytes
- **Osteon**- a central canal with its surrounding lamellae, lacuna, osteocytes.and canaliculi.


Medullary cavity
Periosteum
External circumferential lamellae
Central canal
Osteon
Perforating fibers
Blood vessels
Perforating canals
Central canal
Trabeculae of spongy bone
Internal circumferential lamellae
Central canal
Arteriole
Venule
Nerve
Canaliculi
Osteocyte
Lacuna
Canaliculi

# Haversian System



# Haversian System (Transverse Section)

# Lacunae and Canaliculi



# Cytoplasmic Processes of Osteocytes in Canaliculi



# Bone Histology (cont.)

Spongy (Cancellous) Bone

- consist of an irregular latticework of thin plates of bone called **trabeculae**
- bone marrow located between trabeculae in some bones.
- Lacunae lie within trabeculae where the osteocytes are nourished directly by the blood circulating in the bone marrow
- spongy bone makes up most of the tissue of short, flat, and irregularly shaped bones and most of the epiphyses of long bones.
  - **Hematopoiesis**(blood production) occurs in the spongy bone of hip, ribs, sternum,vertebrae, skull, and ends of some long bones


Proximal epiphysis
Spongy bone
Compact bone
Nutrient foramen
Nutrient vessel
Medullary cavity
Diaphysis
Perforating fibers
Periosteum
Endosteum
Epiphyseal line
Distal epiphysis
Articular cartilage
Coxa

# Spongy bone at the epiphyseal plate



# Bone Formation

- **Ossification**- the process by which bone is formed
  - ossification of embryonic fibrous tissue frame begins around the sixth or seventh week of life and continues throughout adulthood.

Two methods of bone formation

- **intramembranous ossification**- the formation of bone directly on or within fibrous connective tissue membranes.
  - produces the flat bones of the skull and most of the clavicle (collarbone)
- **endochondral ossification**- the formation of bone within a cartilage model.

The two methods do not lead to different structure of bone.

Both methods involve the replacement of a preexisting connective tissue with bone.

# Bone Formation (cont.)

Intramembranous Ossification

flat bones of the skull, mandible (lower jaw bone), and the clavicles (collar bone).

- at the site where bone is to be developed, cells in the mesenchyme began to differentiate into **osteoprogenitor cells** and then to **osteoblasts**. The osteoblast secrete the bony matrix (**osteoid**-soft collagenous tissue void of minerals) until it completely surrounds them. This primary site of bone development is called the **center of ossification.**



# Bone Formation (cont.)

Intramembranous Ossification (cont.)

- when the secretion of the bony matrix completely surrounds the osteoblast, its name changes to **osteocyte** and the small cavity in which it is contained is called a **lacuna** (lacunae, plural).
- the osteocyte extends small cytoplasmic processes outward into the bony matrix forming small channels called **canaliculi**. Soon thereafter, calcium and other mineral salts are deposited and the matrix hardens (**calcification**).



# Bone Formation (cont.)

Intramembranous Ossification (cont.)

- **spongy bone** (the lattice like appearance) is formed when the minute spikes of bone tissue, called **trabeculae**, fuses with one another.
- the spaces between the trabeculae are filled with **marrow** (soft connective tissue that produces red blood cells). On the outside of the bone, the mesenchyme condenses to form the periosteum.




Proximal epiphysis
Spongy bone
Compact bone
Nutrient foramen
Nutrient vessel
Medullary cavity
Diaphysis
Perforating fibers
Periosteum
Endosteum
Epiphyseal line
Distal epiphysis
Articular cartilage
Coxa

# Spongy bone at the epiphyseal plate



# Bone Formation (cont.)

Intramembranous Ossification (cont.)

- eventually, most of the surface layers of the spongy bone are replaced by compact bone, but spongy bone remains in the center of the bone. Much of this newly formed bone will be continually remodeled until final adult size and shaped is reached.


1
Osteoblast
Mesenchymal cell
Collagen fiber
Blood capillary
Center of ossification
2
Osteoblast
Newly calcified bone matrix
Osteocyte in lacuna
Canaliculus
3
Osteoblast
Blood vessel
Mesenchyme condenses
Trabeculae
4
Compact bone
Spongy bone
Periosteum



The process of intramembranous ossification is well seen in histological preparations of the embryonic calvaria. The newly formed bone matrix of developing trabeculae is stained acidophilic (pink) after regular staining. A layer of osteoblasts is present on the surface of the developing trabeculae, whereas osteocytes occupy lacunae in the bone matrix. Even at this early stage osteoclasts are present on the surface of the trabeculae and are active in bone resorption. Primitive blood vessels are seen in the connective tissue located between the trabeculae. At a later stage the connective tissue surrounding the developing flat bone forms the **periosteum**.

# Bone Formation (cont.)

Endochondral Ossification

The replacement of cartilage by bone

Most bones are formed by this process

- Development of the cartilage model
  - cells in the mesenchyme come together in the shape of the future bone
  - the cells differentiate into cartilage producing cells that change the model into hyaline cartilage
  - a membrane called the **perichondrium**(dense regular connective tissue) develops around the cartilage



# Bone Formation (cont.)

Endochondral Ossification (cont.)

- Growth of the cartilage model
  - cartilage model grows in thickness and length.
  - cartilage cells in the midregion trigger calcification leading eventually to death of surrounding cartilage cells due to the inability of nutrients to diffuse though the calcified matrix.
  - the partitions between the lacunae containing the dead cells break down, forming small cavities that will eventually fill with marrow.
  - nutrient arteries penetrate the bone at the midregion through the nutrient foramen.
  - This blood supply stimulate osteoprogenitor cells in the perichondrium to develop into osteoblasts which began to produce a thin layer of compact bone under the perichondrium, known as the **periosteum**.



# Bone Formation (cont.)

Endochondral Ossification (cont.)

- Development of the primary ossification center
  - blood supply continues to growth further into the disintegrating calcified cartilage to stimulate the growth of the primary ossification center, a region where the bone tissue will completely replace most of the cartilage model.
  - in this area osteoblast begin to deposit bone matrix over the remnants of calcified cartilage, forming spongy bone trabeculae. This area, called the **medullary cavity**, is filled by bone marrow.



# Bone Formation (cont.)

Endochondral Ossification (cont.)

- Development of the diaphysis and epiphysis
  - the **diaphysis** (shaft), which was once a solid mass of hyaline cartilage, is replaced by compact bone , the core of which contains a red bone marrow-filled medullary cavity
  - when blood vessels (epiphyseal arteries) enter the **epiphysis**, secondary ossification center develops, usually around the time of birth.



# Bone Formation (cont.)

Endochondral Ossification (cont.)

- Development of the diaphysis and epiphysis
  - in the secondary ossification center, bone formation is similar to that in the primary ossification center, with the following exceptions:
    - spongy bone remains in the interior of the epiphyses (no medullary cavities)
    - hyaline cartilage remains covering the epiphyses as the articular cartilage and between the diaphysis and epiphysis as the **epiphyseal plate** (responsible for the length wise growth of long bones)
  - the epiphyseal plate allows the diaphysis of the bone to <u>increase in length</u> until early childhood
    - rate of growth controlled by hormones , such as **human growth hormone** (hGH)
  - the epiphyseal plate cartilage cells are eventually replaced by bone leading to the cessation of growth. The new structure is called the **epiphyseal line**.
  - <u>growth in diameter</u> occurs as a result of bone destruction by the osteoclast lining the medullary cavity and, at the same time, new bone is being laid down at the outer surface of the bone by the osteoblast of the periosteum.






- Nearest to the epiphysis
- Chondrocytes in a disordered arangement
- Condrocytes are not dividing rapidly and shows no signs or transforming into bone

- New cartilage is produced by interstitial growth
- Multiplying chondrocytes stack up forming longitudinal columns of flattened lacunae

- Chondrocytes mature and enlarge and cease to divide.
- The more mature cells are at the diaphysis end of the column and the less mature ones are at the epiphysis end.

- Thin layer of mineralized matrix
- Death of hypertrophied chondrocytes occurs and the lacunae are invaded by blood vessels
- Osteoblasts from the endosteum, travel with the connective tissue of blood vessels and aggregate on the calcified cartilage surfaces
- New bone matrix is deposited by apositional bone growth, then remodeled

The thickness of the epiphyseal plate remains constant as the length of the diaphysis increases. When the bones reach adult length the plate ossifies and fuses leaving the epiphyseal line. Bones fuse between the age of 12 and 25 and varies between different bones


Endochondral Ossification
Long bone
100 µm

# Bone Homeostasis

Bone growth and maintenance

- Bone is considered never to be at metabolic rest and is constantly being remolded; old bone is being destroyed (osteoclast) and new bone is being laid down (osteoblast).
- a delicate balance is created between the osteoblast and osteoclast.
  - to much new bone tissue causes a condition of bone being abnormally thick and heavy
- This process allows for the removal of worn and injured bone and allow for calcium regulation.

# Osteopetrosis



Bone-within-a-bone appearance of a sagitally sectioned humerus in congenital Osteopetrosis gross. the cone shaped accumulation of bone extending from the metaphyses into the medullary cavity is due to a failure of resorption and remodeling of growing bone that occurs in this condition.

# Bone's Role in Calcium Homeostasis

- Bone is the body's major calcium reservoir, storing 99% of total body calcium.
- Blood calcium is regulated by:
  - Controlling rates of calcium resorption (break down and release into blood)
  - Calcium uptake (deposition from blood into bone

# Calcium Functions in the Body

- Nerve cells depend on having a stable level of calcium ions in the extracellular fluid
- Many enzymes require calcium as a cofactor for enzymatic action
- Blood clotting requires Calcium

Examples of problems related to irregular calcium values (nl 9-11 mg/100ml)

  - Hypercalcemia can cause cardiac arrest
  - Hypocalcemia can cause respiratory arrest

# Parathyroid Hormone

- Secreted by parathyroid glands
- Secreted when blood calcium is too low
- Raises blood calcium by four mechanism
  - stimulates the release of osteoclast-stimulating factor by osteoblast causing increased numbers of osteoclast
  - promotes calcium reabsorption by the kidney
  - increases synthesis of vitamin D
  - inhibits collagen synthesis by osteoblast, thus inhibiting bone deposition

# Hormones That Regulate Calcium

- **Calcitonin-** decreases blood calcium
  - Produced by the parafollicular cells (C cells) of the thyroid gland
  - **Inhibits activity of osteoclasts**
  - **increases the number of osteoblast** thus accelerates calcium deposition into bones
  - Speeds up blood calcium uptake by bone

# Calcitonin (cont)

- **Osteoclast inhibition is very important in children** over adults because
  - osteoclast activity contributes to 5g of blood calcium each day
  - activity in adults only contributes .8g of blood calcium per day
- **Calcitonin deficiency in adults is not known to cause disease**
- Useful in **reducing bone loss in pregnant and lactating women and in the condition of osteoporosis**

# Vitamin D (Calcitriol)

- Produced by the sequential action of the skin, liver, and kidneys
- Principal function is to raise the blood calcium concentration by three mechanism
  - increases calcium absorption by the small intestine (magnesium and phosphate as well)
  - increases the number of osteoclast which liberate calcium and phosphate
  - promotes reabsorption of calcium ions by the kidney from the urine
- Also necessary for bone deposition
  - keeps blood calcium high enough to facilitate bone deposition in bone
  - Low level of vit. D results in softness of bones called rickets in children and osteomalacia in adults.



# Other Factors Effecting Bone Homeostasis

- adequate mineral: calcium, phosphorus, and magnesium, fluoride, iron, and manganese
- Proper amounts of vitamins
  - C for collagen synthesis and for differentiation of osteoblasts to osteocytes
  - K and $B_{12}$ for protein synthesis
  - A stimulates activity of osteoblasts
- Exercise that places stress on bones (weight bearing activities)

# Other Factors Effecting Bone Homeostasis (cont)

- Insulinlike growth factors (IGFs)
  - Produced by bony tissue and liver
  - Stimulated by human growth hormone (hGH) from the anterior pituitary
  - Promotes cell division at the epiphyseal plate and in the periosteum
  - Enhances sysnthesis of protiens needed to build new bone
- Thyroid hormones ($T_3$ & $T_4$) and insulin are also needed for bone growth

# Sex Hormones and Bone Growth

- Androgens and estrogens
  - Responsible for increased osteoblast activity and synthesis of bone matrix
  - Responsible for the sudden growth spurt that occurs during teenage years
  - Estrogen responsible for differences in female skeletal development (ex: widening of pelvis)
    - Responsible for shut down of growth plates in both sexes
      - Lengthwise growth of bone typically ends earlier in females than males due to the higher levels of estrogen in females

| Hormone | Effect | Cause |
| --- | --- | --- |
| **PTH (parathyroid hormone)** | **increases plasma Ca** | **increases renal reabsorption of Ca<br>decreases osteoblast activity<br>increases maturation of preosteoclasts to osteoblasts<br>increases osteoclast activity Ca release from bone** |
| **Calcitonin** | **decreases plasma Ca<br>increases mineral content of bone** | **decreases bone resorption** |
| **Vitamin D** | increases plasma Ca<br>provides adequate Ca for bone formation | increases intestinal absorption of Ca<br>increases renal reabsorption of Ca |

| Hormone | Effect | Cause |
| --- | --- | --- |
| **Thyroid hormones**<br>Hyperthyroidism<br>Hypothyroidism<br>Hypothyroidism + T4 | <br>stops bone growth<br>stops bone growth<br>restores bone growth | increases bone resorption<br>decreases bone production<br>increases proliferation of resting chondrocytes increases proteoglycan synthesis; with the IGF's |
| **Sex steroids increase growth of bone, indirect**<br>Testosterone<br>normal levels<br><br>Estrogens<br>ovariectory (low levels)<br>hi levels | <br><br>increase bone growth<br><br><br>decreases bone growth<br><br>decreases bone mass<br>decreases growth | <br><br>requires GH<br>increase maturation<br>chondrocytes & osteoclasts<br><br>decreases production of bone<br>increases maturation |
| | | |

| Hormone | Effect | Cause |
| --- | --- | --- |
| **Glucocorticoids**<br>Hyperadrenal corticolism<br><br>Normal levels | causes osteoporosis<br><br>normal bone growth | decreases bone formation<br>increases bone resorption<br>decreases IGF levels<br>Synergize with IGFs to increase bone growth, increase IGF receptor number, helps to maintain normal IGF levels in serum |
| **Insulin**<br>Normal levels | required for normal growth | direct effect to increase nutrient uptake by chondrocytes & osteoblasts<br>indirect effect of maintaining IGF levels in serum<br>required for activation of vitamin D (1,25-(OH)2D3) by the kidney |
| **Growth Hormone** | direct effects -- no | indirect effect of maintaining IGF expression |


1
Some stimulus (stress) disrupts homeostasis by causing a decrease in
2
Controlled condition
Blood calcium ($Ca^{2+}$) level
3
Receptors
Parathyroid gland cells detect lowered $Ca^{2+}$ concentration
Input
Increased production of cyclic AMP
4
Control center
PTH gene "turned on"
Output
Increased release of PTH
5
Effectors
Osteoclasts increase bone resorption
Kidneys retain $Ca^{2+}$ in blood, excrete phosphate in urine, and produce calcitriol
6
Response
Increase in blood $Ca^{2+}$ level
7
Return to homeostasis when response brings blood $Ca^{2+}$ level (controlled condition) back to normal

# Exercise and Bone

- Mechanical stress(weight bearing and the pull of gravity) and exercise is vital in bone development
  - mechanical stress causes bone to become stronger through increased deposition of minerals salts and production of collagen fibers
  - stress also increase the production of calcitonin, which inhibits bone reabsorption.
- Aging and Bone
  - aging results in loss of calcium from bone (Osteoporosis).
    - in females it parallels the reduction in estrogen, starting after age 30, accelerating around age 40 to 45, and continues until as much as 30% of bone calcium is lost by age 70.
    - calcium loss does not occur until after age 60 in males.
  - aging also causes a decrease in the deposition of the organic portion of bone (collagen), leading to more inorganic than organic matrix. This causes brittle bone and increased susceptibility to fractures.

# Osteoporosis of the vertebra

# Bone Surface Markings

- Depressions and Openings
  - foramen
  - meatus
  - paranasal sinus
  - fossa
- Processes that Form Joints
  - condyle
  - head
  - facet
- Processes to Which Tendons, Ligaments, and Other Connective Tissue Attach
  - tuberosity
  - spinous process
  - trochanter
  - crest




**Foramen**- an opening through which blood vessels, nerves, or ligaments pass.

**gure 7.12** A radiograph of the pelvic girdle and the articulating nora.






**Meatus**- a tube-like passageway running within a bone, such as the external auditory meatus of the temporal bone.

Paranasal **sinus**- an air-filled cavity within a bone connected to the nasal cavity.






**Tubercle**- a small rough rounded eminence.

**Head**- a rouneded projection that forms a joint and is supported on the constricted portion (neck) of a bone.

**Fossa**- a depression in or on a bone.

**Tuberosity**- large, rounded, usually roughened process.

**Condyle**- large knucklelike prominence that forms a joint.

**Greater Trochanter**

**Trochanter**- a large, blunt projection found only on the femur



**Iliac fossa**

**Crest**




**Crest**- a prominent border or ridge

**gure 7.12** A radiograph of the pelvic girdle and the articulating iora.






Facet

**Facet**- a smooth, flat surface

LUMBAR VERTEBRA, LATERAL VIEW




Spinous Process

**Spinous Process**- a sharp slender projection